



# Snogum



PLANT CHARACTERISTICS	Days to relative maturity/GDU	82/1724
	Cob length (cm)	20.0
	Cob length (inches)	7.9
	Cob width (cm)	5.2
	Cob width (inches)	2.0
	Row number	16
	Plant height (low, medium, tall)	M
	Cob height (low, medium, tall)	M

DISEASE REISISTANCE	Rust common (PS) AVIR (+D)	IR
	Rust common (PS) D-VIR	IR
	Rust common (PS) G-VIR (+D)	IR
	Northern Corn Leaf Blight (ET)	IR
	Stewarts Wilt (PST)	IR
	Maize Dwarf mosaic virus (MDMV)	SU
	Southern Leaf Blight	HR
	Gosses Wilt	

HERBICIDE TOLERANCE	Reaction to Callisto (Mesotrione)	
	Reaction to Accent (Nicosulphuron)	
	Reaction to Laudis (Tembotione)	

KEY	HR	<b>High / Standard Resistance:</b> Describes plant varieties that restrict the growth and development of the specified pest or pathogen under normal pressure when compared to susceptible varieties. HR varieties may exhibit some symptoms or damage under heavy pest or pathogen pressure.
	IR	<b>Intermediate / Moderate Resistance:</b> Describes plant varieties that restrict the growth and development of the specified pest or pathogen but may exhibit a greater range of symptoms or damage compared to HR varieties. IR varieties will still show less severe symptoms or damage than susceptible plan varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
	SU	Susceptibility is the inability of a variety to restrict the growth and development of a specified pest or pathogen.
	*	Means we have insufficient data.

## FIELD PERFORMANCE

Main season. Moderate vigor. High yield, uniform ears with good tip fill on a strong stocky plant. Good disease resistance. Wide harvest window.

## PROCESSING PERFORMANCE

Suits freezing and canning. Cut kernel, Cobbett and whole cob processing. Good white processed color with good eating quality. Excellent uniform cylinder of medium size ears with very good tip fill. Runs well through plants with high recovery.

## MANAGEMENT POINT

Very good holding ability. Good stalk strength. Wide harvest window

## MATURITY SLOT

Plant throughout the season. Widely adaptable outside of MDMV pressure.

Chemical Tolerance is based on findings by the Department of Crop Sciences University of Illinois, USA where Tol = a rating between no or limited apparent injury from the application of the herbicide. Tolerance is defined as the ability of a plant variety to endure abiotic stress without serious consequences for growth, appearance or yield. A tolerant plant will usually show fewer symptoms than sensitive plants when grown under similar conditions of abiotic stress.

Disease ratings are as defined by ISF Position Paper May 2017. Refer to [www.worldseed.org](http://www.worldseed.org) for more information.

The information contained herein is intended as a guide only. Data is based on averages collected from around the world and are indicative only. Varieties should always be trialed in the area they are proposed to be grown. Varietal performance is influenced by many variables, including soil and climatic conditions, cultural and management practices. No liability will be accepted by Snowy River Seeds or its representative for the accuracy of this information.

Notes regarding Rust strains: Common rust races identified in 2001 in USA were not controlled by any of the single genes Rp1-d, Rp1-g or Rp1-i. The effectiveness of rust genes in sweet corn will be determined by the variation of common rust races in each growing area.

Seed is provided with our Standard Seed Treatment or Thiram®, Maxim XL® and Fungaflo® unless otherwise specified.