Sweet Corn Variety Information

Supersweet Yellow (sh2)

Astronaut



PLANT CHARACTERISTICS	Days to relative maturity/GDU	85/1790
	Cob length (cm)	18.6
	Cob length (inches)	7.3
	Cob width (cm)	5.4
	Cob width (inches)	2.2
	Row number	18
	Plant height (low, medium, tall)	М
	Cob height (low, medium, tall)	М
DISEASE REISISTANCE	Rust common (PS) AVIR (+D)	IR
	Rust common (PS) D-VIR	SU
	Rust common (PS) G-VIR (+D)	IR
	Northern Corn Leaf Blight (ET)	IR
	Stewarts Wilt (PST)	
	Maize Dwarf mosaic virus (MDMV)	SU
	Southern Leaf Blight	IR
	Gosses Wilt	
HERBICIDE TOLERANCE	Reaction to Callisto (Mesotrione)	TOL
	Reaction to Accent (Nicosulphuron)	TOL
	Reaction to Laudis (Tembotione)	

HR	High / Standard Resistance: Describes plant varieties that restrict the growth and development of the specified pest or pathogen under normal pressure when compared to susceptible varieties. HR varieties may exhibit some symptoms or damage under heavy pest or pathogen pressure.	
IR	Intermediate / Moderate Resistance: Describes plant varieties that restrict the growth and development of the specified pest or pathogen but may exhibit a greater range of symptoms or damage compared to HR varieties. IR varieties will still show less severe symptoms or damage than susceptible plan varieties when grown under similar environmental conditions and/or pest or pathogen pressure.	
SU	Susceptibility is the inability of a variety to restrict the growth and development of a specified pest or pathogen.	
*	Means we have insufficent data.	

晉



FIELD PERFORMANCE

Full season, full maturity. Medium cob size, excellent tip fill. Stocky plant, nice flags.

PROCESSING PERFORMANCE

Excellent vigour and strong plant, good husk cover, good flags, lovely husk colour, very good fresh market variety.

MANAGEMENT POINT

Good northern corn leaf blight resistance, watch for rust pustules. Will take rust. Can get girthy.

MATURITY SLOT

Plant from the start to the middle of the season, avoid areas where rust can be a problem.

Chemical Tolerance is based on findings by the Department of Crop Sciences University of Illinois, USA where Tol = a rating between no or limited apparent injury from the application of the herbicide. Tolerance is defined as the ability of a plant variety to endure abiotic stress without serious consequences for growth, appearance or yield. A tolerant plant will usually show fewer symptoms than sensitive plants when grown under similar conditions of abiotic stress.

Disease ratings are as defined by ISF Position Paper May 2017. Refer to www.worldseed.org for more information.

The information contained herein is intended as a guide only. Data is based on averages collected from around the world and are indicative only. Varieties should always be trialed in the area they are proposed to be grown. Varietal performance is influenced by many variables, including soil and climatic conditions, cultural and management practices. No liability will be accepted by Snowy River Seeds or its representative for the accuracy of this information.

Notes regarding Rust strains: Common rust races identified in 2001 in USA were not controlled by any of the single genes Rp1-d, Rp1-g or Rp1-i. The effectiveness of rust genes in sweet corn will be determined by the variation of common rust races in each growing area.

Seed is provided with our Standard Seed Treatment or Thiram®, Maxim XL® and Fungaflo® unless otherwise specified.