

Imperator (ZUY7469OY)

PLANT CHARACTERISTICS	Days to relative maturity/GDU	83/1746
	Cob length (cm)	19.8
	Cob length (inches)	7.8
	Cob width (cm)	5.5
	Cob width (inches)	2.2
	Row number	18
	Plant height (low, medium, tall)	М
	Cob height (low, medium, tall)	М
DISEASE REISISTANCE	Rust common (PS) AVIR (+D)	HR
	Rust common (PS) D-VIR	HR
	Rust common (PS) G-VIR (+D)	HR
	Northern Corn Leaf Blight (ET)	IR
	Stewarts Wilt (PST)	
	Maize Dwarf mosaic virus (MDMV)	HR
NCE	Southern Leaf Blight	
""	Gosses Wilt	
ZΞ	Reaction to Callisto (Mesotrione)	
ERBICIDE)LERANCE	Reaction to Accent (Nicosulphuron)	
	Reaction to Laudis (Tembotione)	



FIELD PERFORMANCE

Main season. Excellent vigor with strong cold start ability. High yield and recovery potential. Excellent disease resistance abd good holding ability.

PROCESSING PERFORMANCE

Freezes as both Kernel and Cob. Cans well. Excellent color. Petite kernels. Moderate tip fill. Runs well through processing plants. Late harvest will improve recovery and yield without significant quality loss.

MANAGEMENT POINT

Very good disease resistance with intermediate northern corn leaf blight tolerance. Excellent stalk strength on an attractive plant.

MATURITY SLOT

Can be planted throughout the season.

Chemical Tolerance is based on findings by the Department of Crop Sciences University of Illinois, USA where Tol = a rating between no or limited apparent injury from the application of the herbicide. Tolerance is defined as the ability of a plant variety to endure abiotic stress without serious consequences for growth, appearance or yield. A tolerant plant will usually show fewer symptoms than sensitive plants when grown under similar conditions of abiotic stress.

Disease ratings are as defined by ISF Position Paper May 2017. Refer to www.worldseed.org for more information.

The information contained herein is intended as a guide only. Data is based on averages collected from around the world and are indicative only. Varieties should always be trialed in the area they are proposed to be grown. Varietal performance is influenced by many variables, including soil and climatic conditions, cultural and management practices. No liability will be accepted by Snowy River Seeds or its representative for the accuracy of this information.

Notes regarding Rust strains: Common rust races identified in 2001 in USA were not controlled by any of the single genes Rp1-d, Rp1-g or Rp1-i. The effectiveness of rust genes in sweet corn will be determined by the variation of common rust races in each growing area.

Seed is provided with our Standard Seed Treatment or Thiram®, Maxim XL® and Fungaflo® unless otherwise specified.

HR	when compared to susceptible varieties. HR varieties may exhibit some symptoms or damage under heavy pest or pathogen pressure.
IR	Intermediate / Moderate Resistance: Describes plant varieties that restrict the growth and development of the specified pest or pathogen but may exhibit a greater range of symptoms or damage compared to HR varieties. IR varieties will still show less severe symptoms or damage than susceptible plan varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
SU	Susceptibility is the inability of a variety to restrict the growth and development of a specified pest or pathogen.
*	Means we have insufficent data.

晉

High / Standard Resistance:

Describes plant varieties that restrict the growth and development of the specified